

Dissipative effects from transport and viscous hydrodynamics

Denes Molnar^{1,2} and Pasi Huovinen¹

¹ Purdue University, Physics Department, 525 Northwestern Ave West
Lafayette, IN 47907, USA

² RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY
11973, USA

Abstract. We compare $2 \rightarrow 2$ covariant transport theory and causal Israel-Stewart hydrodynamics in 2+1D longitudinally boost invariant geometry with RHIC-like initial conditions and a conformal $\varepsilon = 3p$ equation of state. The pressure evolution in the center of the collision zone and the final differential elliptic flow $v_2(p_T)$ from the two theories agree remarkably well for a small shear viscosity to entropy density ratio $\eta/s \approx 1/(4\pi)$, and also for a large cross section $\sigma \approx 50$ mb. A key to this agreement is keeping *all* terms in the Israel-Stewart equations of motion. Our results indicate promising prospects for the applicability of Israel-Stewart dissipative hydrodynamics at RHIC, provided the shear viscosity of hot and dense quark-gluon matter is indeed very small for the relevant temperatures $T \sim 200 - 500$ MeV.

PACS numbers: 24.10.Lx, 24.10.Nz, 25.75.Ld

1. Introduction

Recent interest in heavy-ion physics has focused on constraining the transport properties of hot and dense nuclear matter using experimental data from RHIC. A particular open question is the effect of the conjectured “minimal” shear viscosity $\eta = s/(4\pi)$ [1] on the dynamics and observables (s is the entropy density). Studies of dissipation require a suitable theory framework, in principle a non-equilibrium one. However, close to local equilibrium, one can also apply dissipative extensions of ideal (Euler) hydrodynamics [2, 3, 4].

The straightforward Navier-Stokes extension of relativistic ideal hydrodynamics with corrections linear in gradients leads to acausal behaviour and instabilities. An improved formulation proposed by Mueller and later extended by Israel and Stewart [5] (IS) includes second derivatives, which alleviates the causality problem. However, that theory originates from an arbitrary truncation of the entropy current at quadratic order in dissipative corrections (shear and bulk stress, and heat flow), which is not a controlled approximation. Derivations of the IS equations from kinetic theory again rely on an arbitrary truncation of nonequilibrium corrections to the phase space density at quadratic order in momentum (Grad’s 14-moment approximation). In contrast, rigorous (Chapman-Enskog) expansion in small gradients near local equilibrium results in Navier-Stokes theory.

Because of these uncertainties about the region of validity of Israel-Stewart theory, detailed cross-checks against a nonequilibrium approach are paramount. Here we

report on a comparison against covariant transport theory, for conditions expected in $Au + Au$ at $\sqrt{s_{NN}} \sim 200$ GeV at RHIC, and investigate the effect of small shear viscosities on the dynamics and differential elliptic flow $v_2(p_T)$.

2. Covariant transport and Israel-Stewart hydrodynamics

We solve the equations of motion of causal Israel-Stewart dissipative hydrodynamics

$$\partial_\mu T^{\mu\nu} = 0, \quad \partial_\mu N^\mu = 0 \quad (1)$$

$$\begin{aligned} D\pi^{\mu\nu} = & -\frac{1}{\tau_\pi}(\pi^{\mu\nu} - 2\eta\nabla^{\langle\mu}u^{\nu\rangle}) - (u^\mu\pi^{\nu\alpha} + u^\nu\pi^{\alpha\mu})Du_\alpha \\ & - \frac{1}{2}\pi^{\mu\nu}(\partial_\alpha u^\alpha + D\ln\frac{\beta_2}{T}) + 2\pi_\lambda^{\langle\mu}\omega^{\nu\rangle\lambda} \end{aligned} \quad (2)$$

$$T^{\mu\nu} \equiv (\varepsilon + p)u^\mu u^\nu - pg^{\mu\nu} + \pi^{\mu\nu}, \quad N^\mu \equiv nu^\mu \quad (3)$$

in 2+1D longitudinally boost invariant geometry. Here ε , p , u^μ , and n are the local energy density, pressure, flow velocity and particle density; $D \equiv u^\mu\partial_\mu$; the $\langle \rangle$ brackets denote traceless symmetrization and projection orthogonal to the flow

$$A^{\langle\mu\nu\rangle} \equiv \frac{1}{2}\Delta^{\mu\alpha}\Delta^{\nu\beta}(A_{\alpha\beta} + A_{\beta\alpha}) - \frac{1}{3}\Delta^{\mu\nu}\Delta_{\alpha\beta}A^{\alpha\beta}, \quad \Delta^{\mu\nu} \equiv g^{\mu\nu} - u^\mu u^\nu,$$

and $\nabla^\mu \equiv \Delta^{\mu\nu}\partial_\nu$. In the equation for the shear stress $\pi^{\mu\nu}$ we include the term with the vorticity tensor $\omega^{\mu\nu} \equiv (1/2)\Delta^{\mu\alpha}\Delta^{\nu\beta}(\partial_\beta u_\alpha - \partial_\alpha u_\beta)$ that follows from kinetic theory.

The numerical IS solutions are obtained with a modified version of the code used in [7]. Details of the algorithm will be published elsewhere. Covariant transport solutions are calculated via the MPC algorithm [8, 9].

To aid comparison with covariant $2 \rightarrow 2$ transport theory, we take an ideal gas equation of state of massless particles, $\varepsilon = 3p$, and keep particle number conserved. For our conformal system bulk viscosity vanishes, while heat flow is ignored for simplicity. The shear viscosity and shear stress relaxation time of the fluid are matched to the values from kinetic theory [5, 6]: $\eta \approx 6T/(5\sigma)$ and $\tau_\pi \equiv 2\beta_2\eta \approx 9\lambda_{MFP}/5$ (because $\beta_2 = 3/4p$), where an isotropic two-body cross section is utilized for simplicity. We consider two scenarios: i) a constant cross section, and ii) a cross section $\sigma \sim \tau^{2/3}$ growing with proper time $\tau \equiv \sqrt{t^2 - z^2}$. During the initial one-dimensional expansion stage $\eta/s \sim \tau^{2/3}$ grows in the former case, while in the latter case $\eta/s \approx \text{const}$.

Initial conditions expected at RHIC for $Au + Au$ at $\sqrt{s_{NN}} \sim 200$ GeV and impact parameter $b = 8$ fm are modelled through an initial density profile proportional to the (local) number of binary collisions for diffuse gold nuclei, normalized to $dN/d\eta(b=0) = 1000$ to account for the observed $dN_{ch}/d\eta \approx 700$. The system is assumed to start from local thermal equilibrium ($\pi^{\mu\nu} = 0$) at a thermalization time $\tau_0 = 0.6$ fm with uniform initial temperature $T_0 = 385$ MeV.

3. Main results

Figure 1 compares the evolution of the transverse and longitudinal pressure from covariant transport (lines with symbols) and IS hydrodynamics (lines without symbols), averaged over the center of the collision zone $r_T \equiv \sqrt{x^2 + y^2} < 1$ fm for $\eta/s \approx 1/(4\pi)$. At such a low shear viscosity, we find that IS hydrodynamics is a good approximation to covariant transport in the densest region of the collision.




Figure 1. Proper time evolution of the average transverse (T^{xx} , solid) and longitudinal pressure (T^{zz} , dashed) in covariant transport (symbols) and causal Israel-Stewart hydrodynamics (no symbols) near the center of the collision zone ($r_T < 1$ fm) for RHIC-like initial conditions (see text) and $\eta/s \approx 1/(4\pi)$. For clarity, T^{zz} is divided by a factor of 10.

In order to calculate momentum distributions, a freezeout prescription is necessary. Here we apply Cooper-Frye (sudden) freezeout

$$\frac{dN}{dyd^2p_T} = \int d\sigma_\mu(x) p^\mu f(x, \vec{p}) , \quad (4)$$

where f is the phasespace density corresponding to the local fluid variables and $d\sigma_\mu$ is the local normal to the freezeout hypersurface. We choose a constant density hypersurface $n = 0.365 \text{ fm}^{-3}$, which is the density of an ideal gas of massless gluons in chemical equilibrium at $T = 120 \text{ MeV}$ (3 colours).

Figure 2(left) shows elliptic flow from IS hydrodynamics as a function of transverse momentum, for the $\eta/s \approx 1/(4\pi)$ scenario. The dashed curve corresponds to (4) with the local equilibrium ansatz $f_{eq}(x, \vec{p}) = n(x)/[8\pi T(x)^3] \exp[-p^\mu u_\mu/T(x)]$ (we use Boltzmann statistics for consistent comparison with transport). This incorporates dissipative corrections to the evolution of the flow field, temperature and particle number density, but ignores nonequilibrium distortions of the local momentum distributions (i.e., assumes $\pi^{\mu\nu} = 0$). The solid curve is the result for

$$f(x, \vec{p}) = f_{eq}(x, \vec{p}) \left[1 + \frac{p^\mu p^\nu \pi_{\mu\nu}(x)}{8n(x)T^3(x)} \right] \quad (5)$$

that properly takes the local shear stress into account. We find that relative to ideal hydrodynamics (dotted curve), dissipation reduces elliptic flow by $\sim 30\%$, even at such a low $\eta/s \approx 1/(4\pi)$. This reinforces an earlier kinetic theory estimate [10]. Unlike [4], we find that at least one third of the reduction comes from corrections to the ideal hydrodynamic variables (the lower the p_T , the larger the fraction), while the remaining up to two-thirds come from shear stress altering the local momentum distributions.

Finally, we compare in Figure 2(right) differential elliptic flow $v_2(p_T)$ between covariant transport (squares) and IS hydrodynamics (dashed and solid). The dotted line is the ideal hydrodynamics reference. For $\sigma_{gg \rightarrow gg} \approx 47 \text{ mb}$ as in [9] we find excellent agreement between transport and IS hydro (open squares vs dashed line). We also find good agreement for $\eta/s \approx 1/(4\pi)$ (filled squares vs solid line), for which




Figure 2. *Left:* Elliptic flow as a function of p_T for $Au + Au$ at $\sqrt{s_{NN}} \sim 200$ GeV and $b = 8$ fm from ideal hydrodynamics (dotted) and IS hydrodynamics with Cooper-Frye freezeout ignoring (dashed) or incorporating (solid) dissipative corrections to the local momentum distributions (see text) for $\eta/s \approx 1/(4\pi)$. *Right:* Comparison of $v_2(p_T)$ from covariant transport (squares) and IS hydrodynamics (lines) for $\eta/s \approx 1/(4\pi)$ (open squares vs dashed), and $\sigma_{gg \rightarrow gg} \approx 47$ mb (filled squares vs solid). The ideal hydro reference is also shown (dotted).

IS hydro somewhat underpredicts the transport results at higher $p_T > 1.5$ GeV. We caution that the hydro results at high p_T are also sensitive to the freezeout prescription, which deserves further investigation.

The modest $\sim 30\%$ dissipative corrections found agree well with [2] and are much smaller than the effect in [3]. We suspect that this is because [3] ignored the first term in the second line of (2). If the system is near *global* equilibrium, that is appropriate. On the other hand, RHIC-like initial conditions have large gradients, and our comparison to kinetic theory shows that in that case those terms are important.

4. Conclusions

See abstract.

Acknowledgements. We thank RIKEN, Brookhaven National Laboratory and the US Department of Energy [DE-AC02-98CH10886] for providing facilities essential for the completion of this work.

References

- [1] P Kovtun, D T Son and A O Starinets 2005, Phys. Rev. Lett. **94**, 111601
- [2] P Romatschke U Romatschke 2007, Phys. Rev. Lett. **99**, 172301
- [3] H Song and U W Heinz 2008, Phys. Lett. B **658**, 279
- [4] K Dusling and D Teaney 2008, Phys. Rev. C **77**, 034905
- [5] W Israel, 1976, Ann. Phys. **100**, 310; W Israel, J M Stewart 1979, Ann. Phys. **118**, 349
- [6] S R de Groot, W A van Leeuwen, Ch G van Weert 1980, *Relativistic kinetic theory - Principles and applications*, Chs. V, VI.
- [7] P Huovinen, 2005 Nucl. Phys. A **761**, 296
- [8] D Molnar and M Gyulassy 2000, Phys. Rev. C **62**, 054907
- [9] D Molnar and P Huovinen 2005, Phys. Rev. Lett. **94**, 012302

